

FIGURE 1A




FIGURE 1B




FIGURE 1C




FIGURE 2




FIG. 3-2




FIG. 3-3




FIG. 3-4




FIG. 3-5






FIG. 3-6





FIG. 3-7




FIG. 3-8

$$L = \int_0^T \sum_{ik} \left\{ \left( X_{ik} - \hat{X}_{ik} \right)^2 + \mu_{ik} \left[ \hat{X}_{ik} - f \left( \hat{X}_{ik} \right) \right] \right\} dt + K \sum_j P_j^2$$

810
820

actual
estimated

Obtain optimized parameters

- fit patient data
- train on entire recovery period
- minimize error term  $L$
- gradient descent on parameters

$L$	=	Error term
$k$	=	patients
$i$	=	symptoms
$j$	=	parameter
$x$	=	data

FIG. 3-9A




FIG. 3-9B




FIG. 3-10A




FIG. 3-10B




FIG. 3-10C






FIG. 3-10D




FIG. 3-11A




FIG. 3-11B




FIG. 3-12




FIG. 3-13





FIG. 3-14A



(a) Immediate Treatment Effects

FIG. 3-14B




FIG. 3-15






FIG. 3-16




FIG. 3-17





FIG. 4-1




FIG. 4-2




FIGURE 1A




FIGURE 1B




Table 4.1

Prediction of Outcome from Baseline Symptoms														
Symptom	Treatment													
HDRS Item	Other	Tricyclic							MAOI		SSRI	Psychotherapy		ECT
		Ami	Nor	IMI	DMI	Clo	Lev	Map	Phe	Tran	Flu	IPT	CBT	ECT
Severity	5,9-	10-		10-, 11, 16+, 17+	1,8	8,13	3-	3-, 6+	17+		1	16+	7	14-
Depressed Mood	2	10-, 15	4	10-, 11	1,8	8			4	18+	1			14-
Feelings of Guilt		15		11	1,8	8				18	1			14
Suicide	5	15		11	1,8	8				18	1			14
Insomnia: early	2	15	4	11	1,8	8			12,4	18	1			14
Insomnia: middle	2	15-	4	11	1,8	8			12,4	18-	1			14
Insomnia: late	2	15	4	11, 12+	1,8	8			4	18-	1			14
Work and Interests		15	4	11, 12	1,8	8			11	18	1			14-
Retardation	2	10, 15	4	10, 11, 12+	1,8	8			4	18+	1			18
Agitation	2	10, 15		10, 11, 12	1,8	8			12,4	18	1			14-
Anxiety : somatic	2	10, 15	4	10, 11	1,8	8			4	18	1			
Anxiety : psychic	2	10, 15	4	10, 11	1,8	8			4	18	1			
Somatic: gastrointestinal		15-		11	1,8	8				18	1			14-
Somatic: general		10, 15	4	10, 11	1,8	8			4	18	1			
Genital symptoms		15		11, 12	1,8	8			11	18	1			14
Hypochondriasis		15-	4	11	1,8	8			4	18	1			14
Loss of Weight (or change)	2	15	4	11, 12	1,8	8			12,4	18+	1			14
Insight		15+		11	1,8	8				18	1			
Diurnal variation		15	4	11, 12	1,8	8			12,16	18	1			14
Depersonalization and derealization		15		11	1,8	8				18	1			
Paranoid symptoms	2	15		11	1,8	8				18	1			
Obsessive and compulsive	2	15		11	1,8	8				18	1			

## Table 4.2

1. Bowden, C., Schatzberg, A., Rosenbaum, A., Contreras, S., Samson, J., & Dessain, E. (1993). Fluoxetine and desipramine in major depressive disorder. *Journal of Clinical Psychopharmacology*, 13, 305-311.
2. Carney, M., Reynolds, E., & Sheffield, B. (1986). Prediction of outcome in depressive illnesses by the New-castle diagnosis scale: Its relationship to with the unipolar/bipolar and DSM-III systems. *British Journal of Psychiatry*.
3. Filip, V., Hoschl, C., Karen, P., Seifertova, D., Bornova, L., Praskova, H., & Prasko, J. (1993). Predicting therapeutic results with levoprotiline and maprotiline in major depression: The role of the outcome criteria. *British Journal of Psychiatry*, 163, 35-38.
4. Georgotas, A. & McCue, R. E., Cooper, T., Chang, I., Mir, P., & Welkowitz, J. (1987). Clinical Predictors of Response to Antidepressants in Elderly Patients. *Biological Psychiatry*, 22, 733-740.
5. Hinrichsen, G. A. & Hernandez, N. A. (1993). Factors associated with recovery from and relapse into major depressive disorder in the elderly. *American Journal of Psychiatry*, 150, 1820-18 25.
6. Hoencamp, E., Haffmans, P. M. J., Duivenwoorden, H., knegtering, H., & Dijken, W. A. (1994). Predictors of (non-)response in depressed outpatients treated with a three-phase sequential mediation strategy. *Journal of Affective Disorders*, 31, 235-246.
7. Johnson, S. L., Monroe, S., Simons, A., & Thase, M. E. (1994). Clinical characteristics associated with interpersonal depression: symptoms, course, and treatment response. *Journal of Affective Disorders*, 31, 97-109.
8. Joyce, P. R., Mulder, R. T., & Cloninger, C. R. (1994). Temperament predicts clomipramine and desipramine response in major depression. *Journal of Affective Disorders*, 30, 35-46.
9. Katon, W., Lin, E., Von Korff, M., Bush, T., Walker, E., Simon, G., & Robinson, P. (1994). The predictors of depression in primary care. *Journal of Affective Disorders*, 31, 81-90.
10. Katz, M., Koslow, S., Maas, J., Frazer, A., Bowden, C., Casper, R., Croughan, J., Kocsis, J., & Redmond Jr., E. (1987). The timing, specificity, and clinical predictors of tricyclic drug effects in depression. *Psychological Medicine*, 17, 297-309.
11. Kocsis, J. H., Mason, B. J., Frances, A. J., Sweeney, J., Mann, J. J., & Marin, D. (1989). Prediction of response of chronic depression to imipramine. *Journal of Affective Disorders*, 17, 225-260.
12. McGrath, P. J., Stewart, J. W., Harrison, W. M., Ocepek-Welikson, K., Rabkin, J. G., Nunes, E. N., Wagner, S. G., Tricamo, E., Quitkin, F. M., & Klein, D. F. (1992). Predictive value of symptoms of atypical depression for differential treatment outcome. *Journal of Clinical Psycho pharmacology*, 12, 197-202.
13. Nagayama, H., Nagano, K., Ikezaki, A., & Tashiro, T. (1991). Prediction of efficacy of antidepressant by 1-week test therapy in depression. *Journal of Affective Disorders*, 23, 213-216.
14. Pande, A., Krugler, T., Haskett, R., Greden, J., & Grunhaus, L. (1988). Predictors of response to electroconvulsive therapy in major depressive disorder. *Biological Psychiatry*, 24, 91-93.
15. Sauer, H., Kick, H., Minne, H. W., & Schneider, B. (1986). Prediction of the amitriptyline response: Psychopathology versus neuroendocrinology. *International Clinical Psychopharmacology*, 1, 284-295.
16. Sotsky, S. M., Glass, D. R., Shea, T. M., Pilkonis, P. A., Collins, J. F., Elkin, I., Watkins, J. T., Imber, S. D., Leber, W. R., Moyer, J., & Oliveri, M. E. (1991). Patient predictors of response to psychotherapy and pharmacotherapy: Findings in the NIMH treatment of depression collaborative research program. *American Journal of Psychiatry*, 148, 997-1008.
17. Vallejo, J., Gasto, C., Catalan, R., Bulbena, A., & Menchon, J. (1991). Predictors of antidepressant treatment outcome in melancholia: psychosocial, clinical, and biological indicators. *Journal of Affective Disorders*, 21, 151-162.
18. White, K. & White, J. (1986). Tranylcypromine: Patterns and predictors of response. *Journal of Clinical Psychiatry*, 47, 380-382.



FIGURE 1C




FIGURE 2




FIG. 3-2




FIG. 3-3




FIG. 3-4




FIG. 3-5




FIG. 3-6




FIG. 3-7






FIG. 3-8

$$L = \int_0^T \sum_{ik} \left\{ \left( X_{ik} - \hat{X}_{ik} \right)^2 + \mu_{ik} \left( \hat{X}_{ik} - f \left( \hat{X}_{ik} \right) \right) \right\} dt + K \sum_j P_j^2$$

810
820

actual
estimated

Obtain optimized parameters

- fit patient data
- train on entire recovery period
- minimize error term  $L$
- gradient descent on parameters

$L$	=	Error term
$k$	=	patients
$i$	=	symptoms
$j$	=	parameter
$x$	=	data

FIG. 3-9A




FIG. 3-9B




FIG. 3-10A




FIG. 3-10B











FIG. 3-10C





FIG. 3-10D








FIG. 3-11B




FIG. 3-12




FIG. 3-13





FIG. 3-14A



(a) Immediate Treatment Effects




FIG. 3-14B




FIG. 3-15




FIG. 3-16




FIG. 3-17






FIG. 4-1




FIG. 4-2




Table 4.1

Prediction of Outcome from Baseline Symptoms														
Symptom	Treatment													
HDRS Item	Other	Tricyclic							MAOI		SSRI	Psychotherapy		ECT
		Ami	Nor	IMI	DMI	Clo	Lev	Map	Phe	Tran	Flu	IPT	CBT	ECT
Severity	5,9 <del>-</del>	10 <del>-</del>		10 <del>-</del> ,11,16+,17+	1,8	8,13	3 <del>-</del>	3 <del>-</del> ,6+	17+		1	16+	7	14 <del>-</del>
Depressed Mood	2	10 <del>-</del> ,15	4	10 <del>-</del> ,11	1,8	8			4	18+	1			14 <del>-</del>
Feelings of Guilt		15		11	1,8	8				18	1			14
Suicide	5	15		11	1,8	8				18	1			14
Insomnia: early	2	15	4	11	1,8	8			12,4	18	1			14
Insomnia: middle	2	15 <del>-</del>	4	11	1,8	8			12,4	18 <del>-</del>	1			14
Insomnia: late	2	15	4	11,12+	1,8	8			4	18 <del>-</del>	1			14
Work and Interests		15	4	11,12	1,8	8			11	18	1			14 <del>-</del>
Retardation	2	10,15	4	10,11,12+	1,8	8			4	18+	1			18
Agitation	2	10,15		10,11,12	1,8	8			12,4	18	1			14 <del>-</del>
Anxiety : somatic	2	10,15	4	10,11	1,8	8			4	18	1			
Anxiety : psychic	2	10,15	4	10,11	1,8	8			4	18	1			
Somatic: gastrointestinal		15 <del>-</del>		11	1,8	8				18	1			14 <del>-</del>
Somatic: general		10,15	4	10,11	1,8	8			4	18	1			
Genital symptoms		15		11,12	1,8	8			11	18	1			14
Hypochondriasis		15 <del>-</del>	4	11	1,8	8			4	18	1			14
Loss of Weight (or change)	2	15	4	11,12	1,8	8			12,4	18+	1			14
Insight		15+		11	1,8	8				18	1			
Diurnal variation		15	4	11,12	1,8	8			12,16	18	1			14
Depersonalization and derealization		15		11	1,8	8				18	1			
Paranoid symptoms	2	15		11	1,8	8				18	1			
Obsessive, and compulsive	2	15		11	1,8	8				18	1			

## Table 4.2

1. Bowden, C., Schatzberg, A., Rosenbaum, A., Contreras, S., Samson, J., & Dessain, E. (1993). Fluoxetine and desipramine in major depressive disorder. *Journal of Clinical Psychopharmacology*, 13, 305-311.
2. Carney, M., Reynolds, E., & Sheffield, B. (1986). Prediction of outcome in depressive illnesses by the Newcastle diagnosis scale: Its relationship to with the unipolar/bipolar and DSM-III systems. *British Journal of Psychiatry*.
3. Filip, V., Hoschl, C., Karen, P., Seifertova, D., Bornova, L., Praskova, H., & Prasko, J. (1993). Predicting therapeutic results with levoprotiline and maprotiline in major depression: The role of the outcome criteria. *British Journal of Psychiatry*, 163, 35-38.
4. Georgotas, A. & McCue, R. E., Cooper, T., Chang, I., Mir, P., & Welkowitz, J. (1987). Clinical Predictors of Response to Antidepressants in Elderly Patients. *Biological Psychiatry*, 22, 733-740.
5. Hinrichsen, G. A. & Hernandez, N. A. (1993). Factors associated with recovery from and relapse into major depressive disorder in the elderly. *American Journal of Psychiatry*, 150, 1820-18 25.
6. Hoencamp, E., Haffmans, P. M. J., Duivenwoorden, H., knegtering, H., & Dijken, W. A. (1994). Predictors of (non-)response in depressed outpatients treated with a three-phase sequential mediation strategy. *Journal of Affective Disorders*, 31, 235-246.
7. Johnson, S. L., Monroe, S., Simons, A., & Thase, M. E. (1994). Clinical characteristics associated with interpersonal depression: symptoms, course, and treatment response. *Journal of Affective Disorders*, 31, 97-109.
8. Joyce, P. R., Mulder, R. T., & Cloninger, C. R. (1994). Temperament predicts clomipramine and desipramine response in major depression. *Journal of Affective Disorders*, 30, 35-46.
9. Katon, W., Lin, E., Von Korff, M., Bush, T., Walker, E., Simon, G., & Robinson, P. (1994). The predictors of depression in primary care. *Journal of Affective Disorders*, 31, 81-90.
10. Katz, M., Koslow, S., Maas, J., Frazer, A., Bowden, C., Casper, R., Croughan, J., Kocsis, J., & Redmond Jr., E. (1987). The timing, specificity, and clinical predictors of tricyclic drug effects in depression. *Psychological Medicine*, 17, 297-309.
11. Kocsis, J. H., Mason, B. J., Frances, A. J., Sweeney, J., Mann, J. J., & Marin, D. (1989). Prediction of response of chronic depression to imipramine. *Journal of Affective Disorders*, 17, 225-260.
12. McGrath, P. J., Stewart, J. W., Harrison, W. M., Ocepek-Welikson, K., Rabkin, J. G., Nunes, E. N., Wagner, S. G., Tricamo, E., Quitkin, F. M., & Klein, D. F. (1992). Predictive value of symptoms of atypical depression for differential treatment outcome. *Journal of Clinical Psycho pharmacology*, 12, 197-202.
13. Nagayama, H., Nagano, K., Ikezaki, A., & Tashiro, T. (1991). Prediction of efficacy of antidepressant by 1-week test therapy in depression. *Journal of Affective Disorders*, 23, 213-216.
14. Pande, A., Krugler, T., Haskett, R., Greden, J., & Grunhaus, L. (1988). Predictors of response to electroconvulsive therapy in major depressive disorder. *Biological Psychiatry*, 24, 91-93.
15. Sauer, H., Kick, H., Minne, H. W., & Schneider, B. (1986). Prediction of the amitriptyline response: Psychopathology versus neuroendocrinology. *International Clinical Psychopharmacology*, 1, 284-295.
16. Sotsky, S. M., Glass, D. R., Shea, T. M., Pilkonis, P. A., Collins, J. F., Elkin, I., Watkins, J. T., Imber, S. D., Leber, W. R., Moyer, J., & Oliveri, M. E. (1991). Patient predictors of response to psychotherapy and pharmacotherapy: Findings in the NIMH treatment of depression collaborative research program. *American Journal of Psychiatry*, 148, 997-1008.
17. Vallejo, J., Gasto, C., Catalan, R., Bulbena, A., & Menchon, J. (1991). Predictors of antidepressant treatment outcome in melancholia: psychosocial, clinical, and biological indicators. *Journal of Affective Disorders*, 21, 151-162.
18. White, K. & White, J. (1986). Tranylcypromine: Patterns and predictors of response. *Journal of Clinical Psychiatry*, 47, 380-382.